

SPECIAL FEATURE – ESSAY REVIEW

FOREST RESILIENCE, TIPPING POINTS AND GLOBAL CHANGE PROCESSES

Forest resilience and tipping points at different spatio-temporal scales: approaches and challenges

Christopher P. O. Reyer^{1*}, Niels Brouwers², Anja Rammig¹, Barry W. Brook³, Jackie Epila⁴, Robert F. Grant⁵, Milena Holmgren⁶, Fanny Langerwisch¹, Sebastian Leuzinger⁷, Wolfgang Lucht^{1,8}, Belinda Medlyn⁹, Marion Pfeifer¹⁰, Jörg Steinkamp^{11,12}, Mark C. Vanderwel¹³, Hans Verbeeck⁴ and Dora M. Vilella¹⁴

¹Potsdam Institute for Climate Impact Research, Telegrafenberg, PO Box 601203, Potsdam 14412, Germany; ²State Centre of Excellence for Climate Change, Woodland and Forest Heath School of Veterinary and Life Sciences, Murdoch University, Murdoch 6150, WA, Australia; ³Environment Institute and School of Earth and Environmental Sciences The University of Adelaide, Adelaide, SA 5005, Australia; ⁴Laboratory of Plant Ecology Faculty of Bioscience Engineering Ghent University, Coupure Links 653, 9000 Gent, Belgium; ⁵Department of Renewable Resources University of Alberta, Edmonton, AB T6G 2E3, Canada; ⁶Resource Ecology Group Wageningen University, P.O. Box 47, 6700 AA Wageningen, The Netherlands; ⁷Applied Ecology New Zealand Auckland University of Technology, 31-33 Symonds Street, Auckland 1142, New Zealand; ⁸Department of Geography Humboldt-Universität zu Berlin, Unter den Linden 6, D-10099, Berlin, Germany; ⁹Department of Biological Science Macquarie University, North Ryde, NSW 2109, Australia; ¹⁰Department of Life Sciences Imperial College London, Silwood Park Campus, Buckhurst Road, Ascot, Berkshire SL5 7PY, UK; ¹¹Biodiversity and Climate Research Centre (BiK-F), Senckenberganlage 25, 60325 Frankfurt am Main, Germany; ¹²Senckenberg Gesellschaft für Naturforschung, Senckenberganlage 25, 60325 Frankfurt am Main, Germany; ¹³Department of Biology University of Florida, Gainesville, FL 32605, USA; and ¹⁴Laboratório de Ciências Ambientais, CBB, Universidade Estadual do Norte Fluminense (UENF), Av. Alberto Lamego 2000, Campos dos Goytacazes, Rio de Janeiro 28013-600, Brazil

Summary

1. Anthropogenic global change compromises forest resilience, with profound impacts to ecosystem functions and services. This synthesis paper reflects on the current understanding of forest resilience and potential tipping points under environmental change and explores challenges to assessing responses using experiments, observations and models.

2. Forests are changing over a wide range of spatio-temporal scales, but it is often unclear whether these changes reduce resilience or represent a tipping point. Tipping points may arise from interactions across scales, as processes such as climate change, land-use change, invasive species or deforestation gradually erode resilience and increase vulnerability to extreme events. Studies covering interactions across different spatio-temporal scales are needed to further our understanding.

3. Combinations of experiments, observations and process-based models could improve our ability to project forest resilience and tipping points under global change. We discuss uncertainties in changing CO₂ concentration and quantifying tree mortality as examples.

4. Synthesis. As forests change at various scales, it is increasingly important to understand whether and how such changes lead to reduced resilience and potential tipping points. Understanding the mechanisms underlying forest resilience and tipping points would help in assessing risks to ecosystems and presents opportunities for ecosystem restoration and sustainable forest management.

Key-words: climate change, CO₂, drought, environmental change, mortality, plant–climate interactions, regime shifts, review, spatio-temporal scales, vulnerability

*Correspondence author: E-mail: reyer@pik-potsdam.de

Introduction

Global environmental changes in, for instance, atmospheric CO₂ concentration, climate, nitrogen deposition and their interactions are affecting forests around the world (Hyvönen *et al.* 2007; Phillips *et al.* 2009; Lindner *et al.* 2010; Dietze & Moorcroft 2011; Brouwers *et al.* 2012; Poulter *et al.* 2013). These changes occur in combination with direct local anthropogenic drivers of forest change such as land conversion and intensive forest management (e.g. Malhi *et al.* 2008; Barona *et al.* 2010; DeFries *et al.* 2010; Gibbs *et al.* 2010; Kurz 2010; Vayreda *et al.* 2012; Bryan *et al.* 2013). The structural (e.g. reduction in biomass or canopy cover) and functional changes (e.g. changes in community structure) in forests in response to these global and local drivers may compromise the capacity of forest ecosystems to recover from natural or human-induced perturbations (Laurance 2004; Lenton *et al.* 2008). Resilience theory suggests that ecosystems exhibit threshold responses to changing environmental drivers and move towards a new regime when a critical threshold is exceeded (Scheffer *et al.* 2001; Scheffer & Carpenter 2003). However, while evidence for the effects of gradual environmental change on forests exists, the theory of regime shifts has rarely been tested in forest ecosystems and evidence for tipping points at which feedbacks cause ecosystems to enter alternative stable states remains sparse (Hirota *et al.* 2011).

In the scope of the special feature on 'Forest Resilience, Tipping Points and Global Change Processes', we reflect here on the current understanding of forest resilience and potential tipping points under environmental change at different spatial and temporal scales, and the challenges to quantify these ecosystem attributes with experiments, observations and models.

Concepts and definitions

We use the term forest resilience in the sense of Scheffer (2009) as 'the ability of a forest to absorb disturbances and re-organize under change to maintain similar functioning and structure'. This definition is wider than what is understood as 'ecological resilience' (*sensu* Holling 1973) to account for the fact that systems rarely recover to exactly the same pre-disturbance conditions (Scheffer 2009). Furthermore, this definition allows resilience to be framed as the rate of recovery after a disturbance and as the maximum disturbance a forest can absorb before switching to a different ecosystem type (Gunderson 2000; Scheffer 2009) – both being important features of resilience in the context of global change. In systems with multiple stable states, reduced resilience can lead to regime shifts (Scheffer *et al.* 2001). Scheffer (2009) defines a regime shift as a 'relatively sharp change from one regime to a contrasting one, where a regime is a dynamic 'state' of a system with its characteristic stochastic fluctuations and/or cycles'. A tipping point describes a threshold in conditions at which a small change in conditions leads to a strong change in the state of a system (cf. Brook *et al.* 2013).

A typical example to illustrate these concepts is the so-called Amazon forest dieback (White, Cannell & Friend 1999; Cox *et al.* 2004; Malhi *et al.* 2008; Rammig *et al.* 2010). Model simulations including changing precipitation and/or ongoing anthropogenic deforestation in the Amazon show reduced evapotranspiration and hence lower amounts of water recycled for regional precipitation (e.g. Cox *et al.* 2004; Sampaio *et al.* 2007; Costa & Pires 2010). Such an altered hydrological cycle may lead to further tree mortality and reduces forest resilience to recover from fire and drought. Repeated occurrences of strong drought/fire years, for example in the context of El-Niño events, could drive the Amazon past a tipping point into a savanna-like state. Sampaio *et al.* (2007) suggested that at 40% deforestation, such a regime shift could occur in the Amazon basin. Hirota *et al.* (2011) predicted that the south-eastern parts of the Amazon are the least resilient and most likely to turn into savanna or a treeless state.

Another example to illustrate resilience and tipping points in forests at a different spatio-temporal scale is the analysis of drought-induced mortality by Camarero *et al.* (2015). They suggest a framework in which already stressed trees with reduced resilience are further affected by drought up to a point where hydraulic failure occurs or carbon reserves are depleted and additional drought stress can lead to widespread tree mortality and a tipping point.

The spatio-temporal scales at which forest resilience and potential tipping points can be assessed are of critical importance in understanding and interpreting ecological processes (Wiens 1989; Levin 1992). While sophisticated concepts and methods for delineating scales have been discussed recently (Angeler *et al.* 2011; Chave 2013; Allen *et al.* 2014; Nash *et al.* 2014; Soranno *et al.* 2014), we here use the following scale classification to group studies that are relevant to address our main objective of understanding forest resilience and tipping points (Table 1). We refer to 'local' scale as the spatial extent from hundreds of m² up to 10 km², 'regional' scale as larger areas spanning >10 km² to water catchment, ecoregion, country or continental level, and the 'global' scale. For temporal scales, we define 'short term' as periods of up to 10 years capturing processes such as tree mortality and 'long term' as decadal to millennial time periods capturing longer-term processes such as species replacement (see Chapin *et al.* 2004; Hughes *et al.* 2013a).

At which temporal and spatial scales can forest resilience and potential tipping points be assessed?

LOCAL SCALE RESPONSES

At the local scale, short-term and long-term responses to environmental change have been well documented in different types of forests (Table 1). For example, Mediterranean ecosystems have been profoundly transformed by long human use and their response is well studied across different continents. The capacity of Mediterranean ecosystems to respond

Table 1. Examples of changes in forests that are possibly related to reduced resilience and tipping points in different types of forest ecosystems across different temporal and spatial scales as defined in the text. Observed examples in bold, projected or theoretical examples in normal font

	Local (<~10 km ²)	Regional (10 km ² continental scale)	Global
Short-term (<~10 years)	Mediterranean: • Drought and heat-induced tree dieback in south-west Australia (Matusick, Ruthrof & Hardy 2012; Bader <i>et al.</i> 2014) • Drought and fire have induced transition from forest and savannas to shrublands in south Portugal (Acácio <i>et al.</i> 2007, 2009; Acácio & Holmgren 2014) • Drought-induced increase in tree mortality in Spain (Camarero <i>et al.</i> 2015) Different forest types • Drought-induced forest die-back around the world (Allen <i>et al.</i> 2010) • Low safety margin of many species world-wide to drought stress (Choat <i>et al.</i> 2012)	Mediterranean • Drought and heat-induced patchy forest dieback and mortality across a Mediterranean climate ecoregion in south-west Australia (Brouwers <i>et al.</i> 2013; Matusick <i>et al.</i> 2013) • Gradual long-term declines in rainfall and increasing temperatures affecting tree health in south-west Australia (Brouwers <i>et al.</i> 2012) Tropical • Resilience of three alternative stable states of tree covers depends on precipitation in Africa, Australia and South America (Hirota <i>et al.</i> 2011) • Large parts of Amazon and Congo forest could be either forest or savanna, depending on fire regime (Staver, Archibald & Levin 2011) Boreal • Bark beetle-induced tree mortality in western North America (Kurz <i>et al.</i> 2008; Raffa <i>et al.</i> 2008) • Drought-induced dieback of aspen ecotone in mid-continental North America (Michaelian <i>et al.</i> 2011) Different forest types • Drought-induced forest dieback around the world (Allen <i>et al.</i> 2010)	• Catastrophic geophysical events with wide-spread biological consequences (e.g. asteroid strike, supervolcanoes, cosmic radiation from a nearby supernova (e.g. Benton & Twitchett 2003))
Long-term (10–1000 years)	Tropical • Drought decreases biomass in a Monodominant Amazon forest (Nascimento <i>et al.</i> 2014) Different forest types • Drought-induced forest dieback around the world (Allen <i>et al.</i> 2010) • Drought-induced mortality (Steinkamp & Hickler 2015)	Mediterranean • Increasing drought stress in southern Europe (van Oijen <i>et al.</i> 2013) Tropical • Climate-change induced dieback of the Amazon (Cox <i>et al.</i> 2004; Malhi <i>et al.</i> 2009 but see also Cox <i>et al.</i> 2013; Huntingford <i>et al.</i> 2013) Boreal • Boreal biome transitions (Scheffer <i>et al.</i> 2012a) Different forest types • Drought-induced forest die-back around the world (Allen <i>et al.</i> 2010)	• State shift of the global biosphere (Barnosky <i>et al.</i> 2012; Brook <i>et al.</i> 2013; Hughes <i>et al.</i> 2013b; Lenton & Williams 2013)

to disturbances strongly interacts with climate condition. Vegetation changes in the driest regions tend to be persistent and sometimes very difficult to reverse because of positive feedbacks that maintain alternative vegetation states (Holmgren & Scheffer 2001; Acácio *et al.* 2007, 2009; Kefi *et al.* 2007). Mediterranean climate regions are likely to become drier and warmer over the decades to come (Stocker *et al.* 2013), which will likely impact on their resilience as is already occurring in south-west Western Australia (e.g. Brouwers *et al.* 2012). Here, forest tree species are increasingly showing severe dieback and mortality in response to extreme climatic conditions (Matusick, Ruthrof & Hardy 2012; Matusick *et al.* 2013; Bader *et al.* 2014), but whether this constitutes a tipping point is unclear.

Allen *et al.* (2010) concluded that in already dry regions, forests are most vulnerable to further drought possibly indicating reduced resilience. It is precisely these dry regions that have become drier during the second half of the last century (Bates *et al.* 2008; Steinkamp & Hickler 2015). Relative to the small spatial extent of forests with a dry climate, many of the observed drought-induced forest mortality events in Allen *et al.* (2010) occurred in drought-prone forests.

REGIONAL-SCALE RESPONSES

At the regional scale, short- and long-term responses to environmental change have been documented in different forest types (Table 1). Allen *et al.* (2010) provide examples of reduced resilience due to drought-induced tree mortality over both short and long periods of time. Their findings are confirmed by a recent risk analysis of ecosystem vulnerability to drought by van Oijen *et al.* (2013), indicating that drought vulnerability is expected to increase in southern Europe.

Inferring resilience at regional scales is challenging. Hirota *et al.* (2011) and Staver, Archibald & Levin (2011) inferred tropical forest resilience from remotely sensed tree cover distribution. They studied tree cover classes over different precipitation levels that suggest instabilities of forest cover. The 'resilience maps' of Hirota *et al.* (2011) show the probability of finding tropical forests, savannas and treeless conditions in a specific location given the mean annual precipitation. These maps highlight which parts of Africa, Australia and South America are more or less likely to persist in a certain vegetation state if disturbed pointing towards possible tipping points. This approach has also been used to infer the resilience of boreal ecosystems (Scheffer *et al.* 2012a). However, Hanan *et al.* (2014) recently pointed out that such tree cover data sets may be unsuitable for the diagnosis of alternative stable states due to their inherent error distribution. Moreover, remotely sensed data are two-dimensional, time-delayed and often not available as time series.

For the Amazon forests, Cox *et al.* (2004) have projected a tipping point based on climate-change-induced rainfall reductions leading to vegetation–atmosphere feedbacks and further drying using coupled carbon cycle–climate models. However, in more recent studies, Cox *et al.* (2013) and Huntingford *et al.* (2013) show that the Amazon is probably more resilient

than projected in the original Cox *et al.* (2004) study. This change in perspective is due to a better understanding of model uncertainties. Uncertainties with regard to parameterization and structural representation of physiological processes such as CO₂ fertilization or responses to elevated temperatures have been highlighted as being more important than uncertainties related to emission scenarios and climate projections (Huntingford *et al.* 2013). However, another study, using an Earth System Model, found projected changes in dry-season length to be the dominant single factor explaining differences in forest dieback projections (Good *et al.* 2013).

GLOBAL-SCALE RESPONSES

Recently, the potential for an anthropogenically driven global tipping point of the biosphere has been discussed (Barnosky *et al.* 2012; Brook *et al.* 2013; Hughes *et al.* 2013b; Lenton & Williams 2013). Barnosky *et al.* (2012) argue that in the past, state shifts of the totality of the global ecosystem have occurred and that a tipping point of the Earth's biosphere is plausible given the strong and enduring alterations of natural ecosystems by human interventions such as land-use change. Brook *et al.* (2013) highlighted three criteria that need to be fulfilled for such a state shift to occur globally: (i) a spatially homogenous response towards changing environmental conditions, (ii) an interconnection of responses and (iii) a spatial homogeneity of drivers of environmental change. They conclude that spatial heterogeneity in drivers of, and responses to, environmental change as well as the lack of continental connectivity is smoothing global-scale responses and will therefore unlikely result in a global tipping point (Brook *et al.* 2013). Hughes *et al.* (2013b) argue that the speed of change does not determine the presence or the absence of a global tipping point but rather the nonlinear relation between the driving force of ecosystem change and the state of the ecosystem. Lenton & Williams (2013) add that it is the existence of possible feedbacks (and the balance between positive and negative feedbacks) between different components of the Earth system that ultimately determines whether a global tipping of the biosphere exists. They also stress that a clear definition of the 'biosphere' is needed to discuss the possibility of a global tipping point. Lenton & Williams (2013) argue that although a strong global ecological response may be perceived as a tipping point of the biosphere, it is not necessarily associated with a tipping point caused by a feedback within the biosphere itself. They agree with Brook *et al.* (2013) in that interactions of smaller-scale tipping points that influence and trigger each other are more likely than a global tipping point (Lenton & Williams 2013). However, Lenton & Williams (2013) also point out that under certain circumstances, a cascade of tipping points may lead to a tipping point with a global reach.

INTERACTIONS ACROSS SCALES

Besides the individual impacts of natural and anthropogenic pressures, it is perhaps primarily the interaction of processes

across scales that determine forest resilience and tipping points (cf. Gunderson & Holling 2002). For example, the interaction of changes in longer-term mean climate with changes in short-term climate variability determines plant responses to climate change (Reyer *et al.* 2013). In regions where recent climate change has had a significant impact, tipping points may occur when extreme climatic events take place. Hence, continuous slow changes in climatic conditions alter the resilience of forests until extreme events or disturbances, such as prolonged drought, push a specific forest ecosystem across a tipping point into a new state. We therefore argue that, in regions where gradual and persistent long-term drying and warming trends are evident, forest resilience and tree health have likely decreased (e.g. van Mantgem *et al.* 2009; Brouwers *et al.* 2012). This decrease may be a consequence of idiosyncratic interactions of increasing physiological stress, altered natural disturbance regimes and comparably rapid climatic changes, which makes forests vulnerable to regime shift in the future. Similar gradual erosion of resilience through land-use change, invasive species and deforestation may increase vulnerability to extreme events pushing forests towards a tipping point.

Interaction across scales may, however, also increase resilience at larger scales. Higgins & Scheiter (2012), for example, show that although potentially large areas across Africa are threatened by regime shifts in vegetation cover, asynchrony in their timing may dampen their effect on the Earth system globally. Thus, although abrupt transitions can occur locally with potentially devastating effects, they may smooth out on larger spatial scales and prevent strong regional- to global-scale tipping points (Higgins & Scheiter 2012). Similarly, it is important to consider that over larger spatial or temporal scales, there is also a larger potential for adaptation of species and communities to changing conditions, which may increase resilience and further dampen the possibility for a tipping point.

What are the challenges for understanding forest resilience and tipping points under global environmental change?

This section focuses on the challenges in capturing different environmental changes in experiments, observations and models to assess the effects of changing environmental conditions on forests and how this limits our understanding of resilience and tipping points (Table 2). There are, in theory, a number of early warning signals when approaching a regime shift. For instance, these include indicators related to the 'critical slowing down' of a system under stress, a flickering in time series of relevant indicators (e.g. population abundance, nutrient flows) or simply approaching a state where the system has not been observed under current conditions [see Scheffer *et al.* (2012b) and Camarero *et al.* (2015)]. However, the ability to predict a tipping point will remain limited due to the inherent stochasticity of the mechanisms involved (Scheffer *et al.* 2012b).

Another challenge for understanding forest resilience and tipping points is that background conditions are changing, for

example CO₂ concentration and temperature. Changes in human-driven pressures, such as alterations of fire regimes, deforestation, management and their interactions, also need to be considered. Consequently, to deepen our understanding of forest resilience and tipping points integrating insights from experiments, observations and modelling is needed. Experiments may be helpful to determine potential thresholds of ecosystem variables that influence tipping point behaviour, particularly when manipulating whole ecosystems (e.g. Nepstad *et al.* 2007; Brando *et al.* 2008). Observations show conditions under which tipping of a forest ecosystem is possible or has already happened (e.g. Leadley *et al.* 2010). Taken together, this understanding of resilience and tipping points may help to improve process-based models.

Process-based models can be used as diagnostic tools to understand system components that are complicated or costly to measure or as predictive tools that allow using different scenarios to project forest responses to changing environmental conditions (e.g. Reyher *et al.* 2014). Improving process-based models is crucial, because they tend to be very complex and often include mechanisms that are still poorly understood, particularly under changing environmental conditions. With regard to the application of such models to assess resilience and detect tipping points, there is the potential for the implementation of observations and experimentally or theoretically derived mechanisms in models. For example, integrating species-specific rooting profiles into a forest model has been shown to affect competition in mixed forests (Reyer *et al.* 2010). Similar implementations in dynamic global vegetation models could help to better explain forest-savanna interactions as a response to changes in extreme precipitation (Kulmatiski & Beard 2013).

Nonetheless, because models are simplified representations of reality, they should always be considered as generators of hypotheses or projectors of possible pathways and therefore further checked against experimental and independent data on system responses. Most current forest and vegetation models were not meant to simulate tipping points and may not be able to do so because of underpinning assumption like the inclusion of equilibrium pools representing an average steady state response and the lack of feedbacks between vegetation and climate. Moreover, most models still do not contain disturbance processes (e.g. pathogens/insects, wind-throw), which are crucial for simulating mortality of trees and abrupt vegetation shifts (Steinkamp & Hickler 2015). A future challenge is to incorporate these processes in forest models to better simulate resilience and tipping points. In a first attempt, 'tipping point' behaviour could be implemented into models by introducing them explicitly, depending on threshold values. However, in that case, the model would depend totally on the parameter or threshold value imposed. The larger challenge is to build models that do not have these explicit thresholds hard-coded, but have tipping point as an emerging property as a result of the combination of the underlying processes included in the model.

Although most forest models are still limited for exploring tipping points and regime shifts, they are usually able to

Table 2. Examples of changing environmental conditions that are relevant for assessing forest resilience and tipping points and challenges in understanding them in different research approaches

Changing conditions	Observations	Experiments	Models
CO ₂	- Long time series available only close to natural CO₂-sources, which usually represent very specialized environments (e.g. Hättenschwiler <i>et al.</i> 1997) - Europe flux tower records are generally not yet long enough to provide long-term average references (Ciais <i>et al.</i> 2005)	- Only small spatial and temporal coverage of experiments, costly, few in mature stands (Norbby <i>et al.</i> 2010; Bader <i>et al.</i> 2013), no tropical forests but new FACE experiment starting in Amazon - CO₂ increased stepwise vs. gradual change in reality	- CO₂ assumptions in models: sink- vs. source-driven models (Cox <i>et al.</i> 2013; Fatichi, Leuzinger & Kömer 2014) - Assessment of CO₂ response requires N availability to be included in model (Grant 2013)
Temperature	Time series are easy to observe, but the problem is mostly the disentanglement by concurrent changes such as precipitation, CO₂ and nitrogen	- Lack of whole tree warming experiments - Mostly small spatial and temporal coverage, typically glasshouse experiments or open-top chambers	- Temperature responses in models are poorly represented - Acclimation of model parameters is poorly represented
Drought	- Drought and heat often act together, and their interactions need to be considered (Allen <i>et al.</i> 2010; Brouwers <i>et al.</i> 2013; Matusick <i>et al.</i> 2013) - Drought impact responses of forest vegetation are highly variable depending on the positioning within the landscape (Brouwers <i>et al.</i> 2013)	- Expensive, challenging and laborious rainfall exclusion experiments (REE) (Nepstad <i>et al.</i> 2007; Phillips <i>et al.</i> 2010; Misson <i>et al.</i> 2011) - Lateral flow of water not completely controlled in REEs (Fisher <i>et al.</i> 2006) - REEs limited because they alter only rainfall, and not temperature, VPD or radiation (Fisher <i>et al.</i> 2007) - Few actual tropical drought experiments exist (Phillips <i>et al.</i> 2009; da Costa <i>et al.</i> 2010)	Mechanisms of drought-induced mortality such as embolism and cavitation not well covered in many models (Choat <i>et al.</i> 2012; Steinkamp & Hickler 2015)

identify indicators of reduced resilience, for example by predicting changes in biogeochemical cycles. A metric developed by Heyder *et al.* (2011) indicates severe ecosystem change highlighting loss of resilience and the potential approach of tipping points (Gerten *et al.* 2013; Warszawski *et al.* 2013). Recently, simpler ecological models have been developed that are explicitly designed to simulate regime shifts (e.g. Staver & Levin 2012; van Nes *et al.* 2013). In the following sections, we take the example of CO₂ effects and mortality as two key challenges for understanding forest resilience and tipping points under changing environmental conditions in process-based forest models.

EFFECTS OF CHANGING CO₂ CONCENTRATIONS

From a plant's perspective, it is not the role of CO₂ in atmospheric forcing that is most important, but its direct effect on photosynthesis and gas exchange. Since rising CO₂ tends to increase productivity and reduce water use, it may increase forest resilience or lead to the greening of non-forested ecosystems (e.g. Higgins & Scheiter 2012). However, there are counter-examples where rising CO₂ has decreased resilience. In some cases, elevated CO₂ has increased plant leaf area, thus increasing the impact of drought when it occurs (Ghannoum & Way 2011). Also, since elevated CO₂ reduces transpiration, leaf temperatures tend to increase, which may increase the impact of extremely high temperatures (Warren, Norby & Wullschlegel 2011) and decrease resilience to frost events (Barker *et al.* 2005). Increased productivity under elevated CO₂ may also increase litter and reduce litter decomposition rates (Hyvönen *et al.* 2007), leading to increased fuel and higher vulnerability to fire.

Although the effects of elevated CO₂ can be observed close to natural sources of CO₂ (Hättenschwiler *et al.* 1997), derived from time series of measured forest functions under the recent increase in CO₂ (e.g. Keenan *et al.* 2013) or inferred from paleoecological data, experiments and models are most commonly used to study how trees and forests respond to CO₂. The logistics involved in experiments manipulating atmospheric CO₂ levels within existing forests are tremendous and explain why, to date, very few of these experiments exist in natural or semi-natural forests (Norby *et al.* 2010; Leuzinger & Bader 2012; Bader *et al.* 2013). Such experiments remain largely artificial, because the atmospheric CO₂ increase has to occur stepwise instead of gradually as in reality. Furthermore, because transpiration is affected through stomatal closure, forests under elevated atmospheric CO₂ affect local climate (e.g. air humidity). These changes, however, do not feed back on the canopy under study, as the surrounding non-treated trees dominate the climate feedbacks (see Leakey 2009). Tipping points are not to be expected via the direct CO₂ response, but rather via the much more important indirect transpiration response (Holtum & Winter 2010; Hartmann 2011). For example, the rainfall amount and pattern will determine when the CO₂-induced water response starts to become important (Leuzinger & Körner 2010; Hovenden, Newton & Wills

2014). This CO₂-induced water response is likely to be the main process that affects resilience and determines whether a forest tips into an alternate state.

In models, CO₂ effects on forest productivity are key uncertainties. Crucial assumptions relate to whether photosynthesis and stomatal conductance are proportional to CO₂ levels and how this translates to carbon sequestration, how leaf-level processes are coupled to the atmosphere, and how drought affects stomatal conductance or how rainfall is intercepted (Medlyn, Duursma & Zeppel 2011). For example, decreasing productivity under climate change may only be observed in those models that are assuming no interaction between changing temperatures and CO₂-concentrations (Medlyn, Duursma & Zeppel 2011; Reyer *et al.* 2014). Models that include a nitrogen cycle and interactions with CO₂ provide a better representation of CO₂ effects on forest productivity (Grant 2013; Zaehle *et al.* 2014).

MORTALITY

Mortality of individual trees does not mean that resilience of the forest is impaired or a tipping point may occur. However, improving our understanding of mortality will enhance our capacity to simulate forest resilience and tipping points. The mechanistic controls over tree mortality remain a major uncertainty in modelling forest resilience to drought-induced mortality (McDowell *et al.* 2011; Adams *et al.* 2013). Many tree species across the globe operate with small 'safety margins' for xylem embolism and cavitation (Choat *et al.* 2012) and vulnerability to drought is a key factor of tree mortality (Allen *et al.* 2010; Table 1). Mortality is particularly challenging to predict because tree death can occur from a variety of causes, including resource limitation, environmental stress or large-scale natural disturbance. These causes are also strongly interdependent, as reduced carbon or water stores impair a tree's capacity to defend against pests, or to withstand drought. Predicting the onset of mortality thus requires an integrated framework for quantifying critical thresholds and feedbacks in tree responses to physiological stress (McDowell *et al.* 2011).

Regional-scale analyses of tree mortality from forest inventory or forest monitoring plot networks offer outstanding opportunities for quantifying mortality patterns. Long-term, plot-based records have been used to reveal increases in mortality rates over the past 50 years both in the western US and Canada (van Mantgem *et al.* 2009; Peng *et al.* 2011). Such observations can help inform and calibrate the mechanisms used to predict mortality in process-based models, or can serve as a basis for hybrid approaches that combine mechanistic and phenomenological elements to derive predictions of mortality and its role in defining forest resilience or tipping points.

COMBINATION OF METHODS

A combination of experiments, observations and models may help to better understand scale-dependent responses (Fig. 1). Leaf-level CO₂-effects are well understood in the laboratory,




Fig. 1. Temporal and spatial scales at which different research approaches (experiments, observations and models) are available to assess forest resilience and potential tipping points. Drought, pests, fire and other processes that structure the forests such as heatwaves or also deforestation can occur across different spatio-temporal scales. Precipitation, temperature and CO₂ changes cover all spatial scales but only long time scales as we refer to them occurring within the context of climate change and not as part of short-term meteorological variability. The figure is adapted from Gunderson & Holling (2002).

but integrating this mechanistic knowledge with experiments and observations of larger-scale CO₂-effects in models in conjunction with other ecophysiological and demographic processes is necessary to paint a more realistic picture of forest responses to elevated CO₂ (Field, Jackson & Mooney 1995).

Advances have also been made in experiments and observations with regard to the mechanisms underpinning forest growth. While there is empirical evidence that net primary production is at least partly sink – rather than source – driven (Körner 2003; Muller *et al.* 2011; Fatichi, Leuzinger & Körner 2014), current modelling approaches still rely on photosynthesis-driven (i.e. source driven) plant growth (e.g. Bonan *et al.* 2003, 2011; Sitch *et al.* 2003, 2008; Krinner *et al.* 2005). This is highly relevant because photosynthesis often correlates with productivity, and there is a danger of obtaining the right answer for the wrong reason. Sensitivity analyses of vegetation models have shown an unrealistic importance of mostly photosynthesis-related parameters (Pappas *et al.* 2013). However, if tipping points are to be simulated accurately, it is imperative that the causal mechanisms in modelling plant growth are correct, which may involve the reconsideration of the core architecture of vegetation models (Fatichi, Leuzinger & Körner 2014).

Besides combinations of experiments, observation and models, the integration of different model types can aid the understanding of forest resilience. For example, Schelhaas *et al.* (in press) combined simulations of a process-based forest growth model by Reyer *et al.* (2014) with species distribution maps from Hanewinkel *et al.* (2013). By doing this, they could assess where tree species would need to be changed by forest managers as a consequence of changing species

distributions as well as changing productivity under climate change to increase forest resilience.

Concluding remarks

This paper shows that forests are changing over a wide range of spatial and temporal scales. There seems to be more evidence for ongoing changes at smaller spatial and shorter temporal scales and in particular areas such as the Mediterranean. But also larger and longer spatio-temporal scales as well as other forest types face strong changes (Table 1). However, in many cases, it is not clear whether these changes reduce resilience or whether a tipping point is about to be reached. Our synthesis indicates that global-scale disruptions of forest ecosystems are unlikely. We argue that studies covering interactions across a range of spatio-temporal scales are needed to further our understanding of forest resilience and tipping points. Changing CO₂ concentration and tree mortality are processes that require an integration of results from experiments, observations and models to improve the ability of process-based models to simulate forest resilience and potential tipping points.

While acknowledging the important role of gradual changes in affecting resilience of forests under global change and the lack of evidence for mechanisms leading to tipping points, we stress that tipping points are important to investigate from a risk assessment perspective. This means that even though a low likelihood is attributed to a specific high-impact event such as a tipping point, it is still important to discuss whether we are willing to risk such events occurring, especially, because reduced resilience and potential tipping points may have a profound effect on ecosystem functions and services. Such an approach requires an assessment of how society values ecosystem functions and services and would benefit from framing forests as social–ecological systems affected by press and pulse dynamics (cf. Collins *et al.* 2011). At the same time, a better understanding of resilience and tipping points may actually highlight opportunities for ecosystem restoration (e.g. Holmgren *et al.* 2013) or sustainable forest management (Puettmann, Coates & Messier 2009; Rist & Moen 2013), which may provide strong co-benefits for improving human livelihoods.

Acknowledgements

We are grateful to the British Ecological Society and the editorial board of the *Journal of Ecology* for supporting the special feature in which this article is included. The reflections of David Gibson, David Coomes and 2 anonymous referees on an earlier version of this paper are greatly appreciated. We are also indebted to the organizers of the INTECOL 2013 conference at which the idea of this special feature originated in the session called ‘Forest Resilience, Tipping Points and Global Change Processes’. The presenters and the audience of this session are acknowledged for the interesting and informative discussions. Marten Scheffer provided invaluable support. C.R. acknowledges a BES Training and Travel Grant (3817), A.R. the EU-FP7 project ‘AMAZALERT’ (Grant agreement no. 282664) and F.L. the BMBF-Funded Project ‘LEGATO’. D.V. is grateful to CNPq for her Post-doctoral scholarship (236831/2012-7) to support her research in UK. M.P. is supported by the European Research Council Project No. 281986. E.J. is grateful for the grants; Parkyn Bursary (4289-5261) through BES, VLIR-UOS and Ghent University CWO fund.

Data accessibility

This synthesis paper does not use original data.

References

- Acácio, V. & Holmgren, M. (2014) Pathways for resilience in Mediterranean cork oak land use systems. *Annals of Forest Science*, **71**, 5–13.
- Acácio, V., Holmgren, M., Jansen, P.A. & Schrotter, O. (2007) Multiple recruitment limitation causes arrested succession in Mediterranean cork oak systems. *Ecosystems*, **10**, 1220–1230.
- Acácio, V., Holmgren, M., Rego, F., Moreira, F. & Mohren, G.M.J. (2009) Are drought and wildfires turning Mediterranean cork oak forests into persistent shrublands? *Agroforestry Systems*, **76**, 389–400.
- Adams, H.D., Williams, A.P., Xu, C., Rauscher, S.A., Jiang, X. & McDowell, N.G. (2013) Empirical and process-based approaches to climate-induced forest mortality models. *Frontiers in Plant Science*, **4**, 438.
- Allen, C.D., Macalady, A.K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M. *et al.* (2010) A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management*, **259**, 660–684.
- Allen, C.R., Angeler, D.G., Garmestani, A.S., Gunderson, L.H. & Holling, C.S. (2014) Panarchy: theory and application. *Ecosystems*, **17**, 578–589.
- Angeler, D.G., Drakare, S. & Johnson, R.K. (2011) Revealing the organization of complex adaptive systems through multivariate time series modeling. *Ecology and Society*, **16**, 5. doi: 10.5751/ES-04175-160305.
- Bader, M.K.F., Leuzinger, S., Keel, S.G., Siegwolf, R.T.W., Hagedorn, F., Schleppi, P. & Körner, C. (2013) Central European hardwood trees in a high-CO₂ future: synthesis of an 8-year forest canopy CO₂ enrichment project. *Journal of Ecology*, **101**, 1509–1519.
- Bader, M.K.F., Ehrenberger, W., Bitter, R., Stevens, J., Miller, B.P., Chopard, J., Rüger, S., Hardy, G.E.S.J., Poot, P., Dixon, K.W., Zimmermann, U. & Veneklaas, E.J. (2014) Spatio-temporal water dynamics in mature *Banksia menziesii* trees during drought. *Physiologia Plantarum*, **52**, 301–315.
- Barker, D.H., Loveys, B.R., Egerton, J.J.G., Gorton, H., Williams, W.E. & Ball, M.C. (2005) CO₂ enrichment predisposes foliage of a eucalypt to freezing injury and reduces spring growth. *Plant, Cell & Environment*, **28**, 1506–1515.
- Barnosky, A.D., Hadly, E.A., Bascompte, J., Berlow, E.L., Brown, J.H., Fortelius, M. *et al.* (2012) Approaching a state shift in Earth's biosphere. *Nature*, **486**, 52–58.
- Barona, E., Ramankutty, N., Hyman, G. & Coomes, O. (2010) The role of pasture and soybean in deforestation of the Brazilian Amazon. *Environmental Research Letters*, **5**, 024002.
- Bates, B.C., Hope, P., Ryan, B., Smith, I. & Charles, S. (2008) Key findings from the Indian Ocean climate initiative and their impact on policy development in Australia. *Climatic Change*, **89**, 339–354.
- Benton, M.J. & Twitchett, R.J. (2003) How to kill (almost) all life: the end-Permian extinction event. *Trends in Ecology & Evolution*, **18**, 358–365.
- Bonan, G.B., Levis, S., Sitch, S., Vertenstein, M. & Oleson, K.W. (2003) A dynamic global vegetation model for use with climate models: concepts and description of simulated vegetation dynamics. *Global Change Biology*, **9**, 1543–1566.
- Bonan, G.B., Lawrence, P.J., Oleson, K.W., Levis, S., Jung, M., Reichstein, M., Lawrence, D.M. & Swenson, S.C. (2011) Improving canopy processes in the Community Land Model version 4 (CLM4) using global flux fields empirically inferred from FLUXNET data. *Journal of Geophysical Research: Biogeosciences*, **116**, G02014.
- Brando, P.M., Nepstad, D.C., Davidson, E.A., Trumbore, S.E., Ray, D. & Camargo, P.n. (2008) Drought effects on litterfall, wood production and belowground carbon cycling in an Amazon forest: results of a throughfall reduction experiment. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **363**, 1839–1848.
- Brook, B.W., Ellis, E.C., Perring, M.P., Mackay, A.W. & Blomqvist, L. (2013) Does the terrestrial biosphere have planetary tipping points? *Trends in Ecology & Evolution*, **28**, 396–401.
- Brouwers, N.C., Mercer, J., Lyons, T., Poot, P., Veneklaas, E. & Hardy, G. (2012) Climate and landscape drivers of tree decline in a Mediterranean ecosystem. *Ecology and Evolution*, **3**, 67–79.
- Brouwers, N.C., Matusick, G., Ruthrof, K., Lyons, T. & Hardy, G. (2013) Landscape-scale assessment of tree crown dieback following extreme drought and heat in a Mediterranean eucalypt forest ecosystem. *Landscape Ecology*, **28**, 69–80.
- Bryan, J.E., Shearman, P.L., Asner, G.P., Knapp, D.E., Aoro, G. & Lokes, B. (2013) Extreme differences in forest degradation in Borneo: comparing practices in Sarawak, Sabah, and Brunei. *PLoS ONE*, **8**, e69679.
- Camarero, J.J., Gazol, A., Sangüesa-Barreda, G., Oliva, J. & Vicente-Serrano, S.M. (2015) To die or not to die: early-warning signals of drought-induced tree die-off. *Journal of Ecology*, **103**, 43–56.
- Chapin, F.S., Callaghan, T.V., Bergeron, Y., Fukuda, M., Johnstone, J.F., Juday, G. & Zimov, S.A. (2004) Global change and the boreal forest: thresholds, shifting states or gradual change? *AMBIO: A Journal of the Human Environment*, **33**, 361–365.
- Chave, J. (2013) The problem of pattern and scale in ecology: what have we learned in 20 years? *Ecology Letters*, **16**, 4–16.
- Choat, B., Jansen, S., Brodribb, T.J., Cochard, H., Delzon, S., Bhaskar, R. *et al.* (2012) Global convergence in the vulnerability of forests to drought. *Nature*, **491**, 752–755.
- Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogee, J., Allard, V. *et al.* (2005) Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature*, **437**, 529–533.
- Collins, S.L., Carpenter, S.R., Swinton, S.M., Orenstein, D.E., Childers, D.L., Gragson, T.L. *et al.* (2011) An integrated conceptual framework for long-term social-ecological research. *Frontiers in Ecology and the Environment*, **9**, 351–357.
- Costa, M.H. & Pires, G.F. (2010) Effects of Amazon and Central Brazil deforestation scenarios on the duration of the dry season in the arc of deforestation. *International Journal of Climatology*, **30**, 1970–1979.
- Cox, P.M., Betts, R.A., Collins, M., Harris, P.P., Huntingford, C. & Jones, C.D. (2004) Amazonian forest dieback under climate-carbon cycle projections for the 21st century. *Theoretical and Applied Climatology*, **78**, 137–156.
- Cox, P.M., Pearson, D., Booth, B.B., Friedlingstein, P., Huntingford, C., Jones, C.D. & Luke, C.M. (2013) Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability. *Nature*, **494**, 341–344.
- DeFries, R.S., Rudel, T., Uriarte, M. & Hansen, M. (2010) Deforestation driven by urban population growth and agricultural trade in the twenty-first century. *Nature Geoscience*, **3**, 178–181.
- Dietze, M.C. & Moorcroft, P.R. (2011) Tree mortality in the eastern and central United States: patterns and drivers. *Global Change Biology*, **17**, 3312–3326.
- Faticchi, S., Leuzinger, S. & Körner, C. (2014) Moving beyond photosynthesis: from carbon source to sink-driven vegetation modeling. *New Phytologist*, **201**, 1086–1095.
- Field, C.B., Jackson, R.B. & Mooney, H.A. (1995) Stomatal responses to increased CO₂: implications from the plant to the global scale. *Plant, Cell and Environment*, **18**, 1214–1225.
- Fisher, R.A., Williams, M., Do Vale, R.L., Da Costa, A.L. & Meir, P. (2006) Evidence from Amazonian forests is consistent with isohydric control of leaf water potential. *Plant, Cell & Environment*, **29**, 151–165.
- Fisher, R.A., Williams, M., Da Costa, A.L., Malhi, Y., Da Costa, R.F., Almeida, S. & Meir, P. (2007) The response of an Eastern Amazonian rain forest to drought stress: results and modelling analyses from a throughfall exclusion experiment. *Global Change Biology*, **13**, 2361–2378.
- Gerten, D., Lucht, W., Ostberg, S., Heinke, J., Kowarsch, M., Kreft, H., Kundzewicz, Z.W., Rastgooy, J., Warren, R. & Schellnhuber, H.J. (2013) Asynchronous exposure to global warming: freshwater resources and terrestrial ecosystems. *Environmental Research Letters*, **8**, 034032.
- Ghannoum, O. & Way, D.A. (2011) On the role of ecological adaptation and geographic distribution in the response of trees to climate change. *Tree Physiology*, **31**, 1273–1276.
- Gibbs, H.K., Ruesch, A.S., Achard, F., Clayton, M.K., Holmgren, P., Ramankutty, N. & Foley, J.A. (2010) Tropical forests were the primary sources of new agricultural land in the 1980s and 1990s. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 16732–16737.
- Good, P., Jones, C., Lowe, J., Betts, R. & Gedney, N. (2013) Comparing tropical forest projections from two generations of hadley centre earth system models, HadGEM2-ES and HadCM3LC. *Journal of Climate*, **26**, 495–511.
- Grant, R.F. (2013) Modelling changes in nitrogen cycling to sustain increases in forest productivity under elevated atmospheric CO₂ and contrasting site conditions. *Biogeosciences*, **10**, 7703–7721.
- Gunderson, L.H. (2000) Ecological resilience – in theory and application. *Annual Review of Ecology and Systematics*, **31**, 425–439.
- Gunderson, L. & Holling, C.S. (eds) (2002) *Panarchy - Understanding Transformations in Human and Natural Systems*. Island Press, Washington DC, USA, pp. 507.

- Hanan, N.P., Tredennick, A.T., Prihodko, L., Bucini, G. & Dohn, J. (2014) Analysis of stable states in global savannas: is the CART pulling the horse? *Global Ecology and Biogeography*, **23**, 259–263.
- Hanewinkel, M., Cullmann, D.A., Schelhaas, M.-J., Nabuurs, G.-J. & Zimmermann, N.E. (2013) Climate change may cause severe loss in the economic value of European forest land. *Nature Climate Change*, **3**, 203–207.
- Hartmann, H. (2011) Will a 385 million year-struggle for light become a struggle for water and for carbon? – How trees may cope with more frequent climate change-type drought events. *Global Change Biology*, **17**, 642–655.
- Hättenschwiler, S., Miglietta, F., Raschi, A. & Körner, C. (1997) Thirty years of *in situ* tree growth under elevated CO₂: a model for future forest responses? *Global Change Biology*, **3**, 463–471.
- Heyder, H., Schaphoff, S., Gerten, D. & Lucht, W. (2011) Risk of severe climate change impact on the terrestrial biosphere. *Environmental Research Letters*, **6**, 034036.
- Higgins, S.I. & Scheiter, S. (2012) Atmospheric CO₂ forces abrupt vegetation shifts locally, but not globally. *Nature*, **488**, 209–212.
- Hirota, M., Holmgren, M., Van Nes, E.H. & Scheffer, M. (2011) Global resilience of tropical forest and savanna to critical transitions. *Science*, **334**, 232–235.
- Holling, C.S. (1973) Resilience and stability of ecological systems. *Annual Review of Ecology and Systematics*, **4**, 1–23.
- Holmgren, M. & Scheffer, M. (2001) El Niño as a window of opportunity for the restoration of degraded arid ecosystems. *Ecosystems*, **4**, 151–159.
- Holmgren, M., Hirota, M., van Nes, E.H. & Scheffer, M. (2013) Effects of interannual climate variability on tropical tree cover. *Nature Climate Change*, **3**, 755–758.
- Holtum, J.A.M. & Winter, K. (2010) Elevated [CO₂] and forest vegetation: more a water issue than a carbon issue? *Functional Plant Biology*, **37**, 694–702.
- Hovenden, M.J., Newton, P.C.D. & Wills, K.E. (2014) Seasonal not annual rainfall determines grassland biomass response to carbon dioxide. *Nature*, **511**, 583–586.
- Hughes, T.P., Linares, C., Dakos, V., van de Leemput, I.A. & van Nes, E.H. (2013a) Living dangerously on borrowed time during slow, unrecognized regime shifts. *Trends in Ecology & Evolution*, **28**, 149–155.
- Hughes, T.P., Carpenter, S., Rockström, J., Scheffer, M. & Walker, B. (2013b) Multiscale regime shifts and planetary boundaries. *Trends in Ecology & Evolution*, **28**, 389–395.
- Huntingford, C., Zelazowski, P., Galbraith, D., Mercado, L.M., Sitch, S., Fisher, R. *et al.* (2013) Simulated resilience of tropical rainforests to CO₂-induced climate change. *Nature Geoscience*, **6**, 268–273.
- Hyvönen, R., Ågren, G.I., Linder, S., Persson, T., Cotrufo, M.F., Ekblad, A. *et al.* (2007) The likely impact of elevated [CO₂], nitrogen deposition, increased temperature and management on carbon sequestration in temperate and boreal forest ecosystems: a literature review. *New Phytologist*, **173**, 463–480.
- Keenan, T.F., Hollinger, D.Y., Bohrer, G., Dragoni, D., Munger, J.W., Schmid, H.P. & Richardson, A.D. (2013) Increase in forest water-use efficiency as atmospheric carbon dioxide concentrations rise. *Nature*, **499**, 324–327.
- Kefi, S., Rietkerk, M., Alados, C.L., Pueyo, Y., Papanastasis, V.P., ElAich, A. & de Ruiter, P.C. (2007) Spatial vegetation patterns and imminent desertification in Mediterranean arid ecosystems. *Nature*, **449**, 213–217.
- Körner, C. (2003) Carbon limitation in trees. *Journal of Ecology*, **91**, 4–17.
- Krinner, G., Viovy, N., de Noblet-Ducoudré, N., Ogée, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S. & Prentice, I.C. (2005) A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system. *Global Biogeochemical Cycles*, **19**, GB1015.
- Kulmatiski, A. & Beard, K.H. (2013) Woody plant encroachment facilitated by increased precipitation intensity. *Nature Climate Change*, **3**, 833–837.
- Kurz, W.A. (2010) An ecosystem context for global gross forest cover loss estimates. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 9025–9026.
- Kurz, W.A., Dymond, C.C., Stinson, G., Rampley, G.J., Neilson, E.T., Carroll, A.L., Ebata, T. & Safranyik, L. (2008) Mountain pine beetle and forest carbon feedback to climate change. *Nature*, **452**, 987–990.
- Laurance, W.F. (2004) Forest-climate interactions in fragmented tropical landscapes. *Philosophical Transactions of the Royal Society B*, **359**, 345–352.
- Leadley, P., Pereira, H.M., Alkamade, R., Fernandez-Manjarrés, J.F., Proença, V., Scharlemann, J.P.W. & Walpole, M.J. (2010) Biodiversity Scenarios: Projections of 21st century change in biodiversity and associated ecosystem services, Montreal.
- Leakey, A.D.B. (2009) Rising atmospheric carbon dioxide concentration and the future of C4 crops for food and fuel. *Proceedings of the Royal Society B*, **276**, 2333–2343.
- Lenton, T.M. & Williams, H.T.P. (2013) On the origin of planetary-scale tipping points. *Trends in Ecology & Evolution*, **28**, 380–382.
- Lenton, T.M., Held, H., Kriegler, E., Hall, J.W., Lucht, W., Rahmstorf, S. & Schellnhuber, H.J. (2008) Tipping elements in the Earth's climate system. *Proceedings of the National Academy of Sciences of the United States of America*, **105**, 1786–1793.
- Leuzinger, S. & Bader, M.K.F. (2012) Experimental vs. modeled water use in mature Norway spruce (*Picea abies*) exposed to elevated CO₂. *Frontiers in Plant Science*, **3**, 229.
- Leuzinger, S. & Körner, C. (2010) Rainfall distribution is the main driver of runoff under future CO₂-concentration in a temperate deciduous forest. *Global Change Biology*, **16**, 246–254.
- Levin, S.A. (1992) The problem of pattern and scale in ecology: the Robert H. MacArthur Award Lecture. *Ecology*, **73**, 1943–1967.
- Lindner, M., Maroschek, M., Netherer, S., Kremer, A., Barbati, A., Garcia-Gonzalo, J., Seidl, R., Delzon, S., Corona, P., Kolström, M., Lexer, M.J. & Marchetti, M. (2010) Climate change impacts, adaptive capacity, and vulnerability of European forest ecosystems. *Forest Ecology and Management*, **259**, 698–709.
- Malhi, Y., Roberts, J.T., Betts, R.A., Killeen, T.J., Li, W. & Nobre, C.A. (2008) Climate change, deforestation, and the fate of the Amazon. *Science*, **319**, 169–172.
- Malhi, Y., Aragão, L.E.O.C., Galbraith, D., Huntingford, C., Fisher, R., Zelazowski, P., Sitch, S., McSweeney, C. & Meir, P. (2009) Exploring the likelihood and mechanism of a climate-change-induced dieback of the Amazon rainforest. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 20610–20615.
- van Mantgem, P.J., Stephenson, N.L., Byrne, J.C., Daniels, L.D., Franklin, J.F., Fulé, P.Z., Harmon, M.E., Larson, A.J., Smith, J.M., Taylor, A.H. & Veblen, T.T. (2009) Widespread increase of tree mortality rates in the Western United States. *Science*, **323**, 521–524.
- Matusick, G., Ruthrof, K.X. & Hardy, G.E.S.J. (2012) Drought and heat triggers sudden and severe dieback in a dominant Mediterranean-type woodland species. *Open Journal of Forestry*, **2**, 183–186.
- Matusick, G., Ruthrof, K.X., Brouwers, N.C., Dell, B. & Hardy, G.J. (2013) Sudden forest canopy collapse corresponding with extreme drought and heat in a mediterranean-type eucalypt forest in southwestern Australia. *European Journal of Forest Research*, **132**, 497–510.
- McDowell, N.G., Beerling, D.J., Breshears, D.D., Fisher, R.A., Raffa, K.F. & Stitt, M. (2011) The interdependence of mechanisms underlying climate-driven vegetation mortality. *Trends in Ecology & Evolution*, **26**, 523–532.
- Medlyn, B.E., Duursma, R.A. & Zeppel, M.J.B. (2011) Forest productivity under climate change: a checklist for evaluating model studies. *Wiley Interdisciplinary Reviews: Climate Change*, **2**, 332–355.
- Michaelian, M., Hogg, E.H., Hall, R.J. & Arseneault, E. (2011) Massive mortality of aspen following severe drought along the southern edge of the Canadian boreal forest. *Global Change Biology*, **17**, 2084–2094.
- Misson, L., Degueldre, D., Collin, C., Rodriguez, R., Rocheteau, A., Ourcival, J.-M. & Rambal, S. (2011) Phenological responses to extreme droughts in a Mediterranean forest. *Global Change Biology*, **17**, 1036–1048.
- Muller, B., Pantin, F., Génard, M., Turc, O., Freixes, S., Piques, M. & Gibon, Y. (2011) Water deficits uncouple growth from photosynthesis, increase C content, and modify the relationships between C and growth in sink organs. *Journal of Experimental Botany*, **62**, 1715–1729.
- Nascimento, M.T., da Silva Carvalho, L.C., Barbosa, R.I. & Villela, D.M. (2014) Variation in floristic composition, demography and above-ground biomass over a 20-year period in an Amazonian monodominant forest. *Plant Ecology & Diversity*, **7**, 293–303.
- Nash, K.L., Allen, C.R., Angeler, D.G., Barichev, C., Eason, T., Garmestani, A.S. *et al.* (2014) Discontinuities, cross-scale patterns, and the organization of ecosystems. *Ecology*, **95**, 654–667.
- Nepstad, D.C., Tohver, I.M., Ray, D., Moutinho, P. & Cardinot, G. (2007) Mortality of large trees and lianas following experimental drought in an Amazon forest. *Ecology*, **88**, 2259–2269.
- van Nes, E.H., Hirota, M., Holmgren, M. & Scheffer, M. (2013) Tipping points in tropical tree cover: linking theory to data. *Global Change Biology*, **20**, 1016–1021.
- Norby, R.J., Warren, J.M., Iversen, C.M., Medlyn, B.E. & McMurtrie, R.E. (2010) CO₂ enhancement of forest productivity constrained by limited nitrogen availability. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 19368–19373.
- van Oijen, M., Beer, C., Cramer, W., Rammig, A., Reichstein, M., Rolinski, S. & Soussana, J.-F. (2013) A novel probabilistic risk analysis to determine the vulnerability of ecosystems to extreme climatic events. *Environmental Research Letters*, **8**, 015032.

- Pappas, C., Fatichi, S., Leuzinger, S., Wolf, A. & Burlando, P. (2013) Sensitivity analysis of a process-based ecosystem model: pinpointing parameterization and structural issues. *Journal of Geophysical Research: Biogeosciences*, **118**, 505–528.
- Peng, C., Ma, Z., Lei, X., Zhu, Q., Chen, H., Wang, W., Liu, S., Li, W., Fang, X. & Zhou, X. (2011) A drought-induced pervasive increase in tree mortality across Canada's boreal forests. *Nature Climate Change*, **1**, 467–471.
- Phillips, O.L., Aragão, L.E.O.C., Lewis, S.L., Fisher, J.B., Lloyd, J., López-González, G. *et al.* (2009) Drought sensitivity of the Amazon rainforest. *Science*, **323**, 1344–1347.
- Phillips, O.L., van der Heijden, G., Lewis, S.L., López-González, G., Aragão, L.E.O.C., Lloyd, J. *et al.* (2010) Drought–mortality relationships for tropical forests. *New Phytologist*, **187**, 631–646.
- Poulter, B., Pederson, N., Liu, H., Zhu, Z., D'Arrigo, R., Ciais, P., Davi, N., Frank, D., Leland, C., Myneni, R., Piao, S. & Wang, T. (2013) Recent trends in Inner Asian forest dynamics to temperature and precipitation indicate high sensitivity to climate change. *Agricultural and Forest Meteorology*, **178–179**, 31–45.
- Puettmann, K., Coates, K.D. & Messier, C. (2009) *A critique of Silviculture - Managing for Complexity* Island Press, Washington, DC.
- Raffa, K.F., Aukema, B.H., Bentz, B.J., Carroll, A.L., Hicke, J.A., Turner, M.G. & Romme, W.H. (2008) Cross-scale drivers of natural disturbances prone to anthropogenic amplification: the dynamics of bark beetle eruptions. *BioScience*, **58**, 501–517.
- Rammig, A., Jupp, T., Thonicke, K., Tietjen, B., Heinke, J., Ostberg, S., Lucht, W., Cramer, W. & Cox, P. (2010) Estimating the risk of Amazonian forest dieback. *New Phytologist*, **187**, 694–706.
- Reyer, C., Lasch, P., Mohren, G.M.J. & Sterck, F.J. (2010) Inter-specific competition in mixed forests of Douglas-fir (*Pseudotsuga menziesii*) and common beech (*Fagus sylvatica*) under climate change - a model-based analysis. *Annals of Forest Science*, **67**, 805.
- Reyer, C., Leuzinger, S., Rammig, A., Wolf, A., Bartholomeus, R.P., Bonfante, A. *et al.* (2013) A plant's perspective of extremes: terrestrial plant responses to changing climatic variability. *Global Change Biology*, **19**, 75–89.
- Reyer, C., Lasch-Born, P., Suckow, F., Gutsch, M., Murawski, A. & Pilz, T. (2014) Projections of regional changes in forest net primary productivity for different tree species in Europe driven by climate change and carbon dioxide. *Annals of Forest Science*, **71**, 211–225.
- Rist, L. & Moen, J. (2013) Sustainability in forest management and a new role for resilience thinking. *Forest Ecology and Management*, **310**, 416–427.
- Sampaio, G., Nobre, C., Costa, M.H., Satyamurty, P., Soares-Filho, B.S. & Cardoso, M. (2007) Regional climate change over eastern Amazonia caused by pasture and soybean cropland expansion. *Geophysical Research Letters*, **34**, L17709.
- Scheffer, M. (2009) *Critical Transitions in Nature and Society*. Princeton University Press, Princeton, USA.
- Scheffer, M. & Carpenter, S.R. (2003) Catastrophic regime shifts in ecosystems: linking theory to observation. *Trends in Ecology & Evolution*, **18**, 648–656.
- Scheffer, M., Carpenter, S., Foley, J.A., Folke, C. & Walker, B. (2001) Catastrophic shifts in ecosystems. *Nature*, **413**, 591–596.
- Scheffer, M., Hirota, M., Holmgren, M., Van Nes, E.H. & Chapin, F.S. (2012a) Thresholds for boreal biome transitions. *Proceedings of the National Academy of Sciences of the United States of America*, **109**, 21384–21389.
- Scheffer, M., Carpenter, S.R., Lenton, T.M., Bascompte, J., Brock, W., Dakos, V., van de Koppel, J., van de Leemput, I.A., Levin, S.A., van Nes, E.H., Pascual, M. & Vandermeer, J. (2012b) Anticipating critical transitions. *Science*, **338**, 344–348.
- Schellhaas, M., Nabuurs, G., Hengeveld, G., Reyer, C., Hanewinkel, M., Zimmermann, N. & Cullmann, D. (in press) Adaptive forest management to account for climate change-induced productivity and species suitability changes in Europe. *Regional Environmental Change*.
- Sitch, S., Smith, B., Prentice, I.C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J.O., Levis, S., Lucht, W., Sykes, M.T., Thonicke, K. & Venevsky, S. (2003) Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Global Change Biology*, **9**, 161–185.
- Sitch, S., Huntingford, C., Gedney, N., Levy, P.E., Lomas, M., Piao, S.L. *et al.* (2008) Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five Dynamic Global Vegetation Models (DGVMs). *Global Change Biology*, **14**, 2015–2039.
- Soranno, P.A., Cheruvelil, K.S., Bissell, E.G., Bremigan, M.T., Downing, J.A., Fergus, C.E. *et al.* (2014) Cross-scale interactions: quantifying multi-scaled cause-effect relationships in macrosystems. *Frontiers in Ecology and the Environment*, **12**, 65–73.
- Staver, A.C., Archibald, S. & Levin, S.A. (2011) The global extent and determinants of savanna and forest as alternative biome states. *Science*, **334**, 230–232.
- Staver, A.C. & Levin, S. (2012) Integrating theoretical climate and fire effects on savanna and forest systems. *The American Naturalist*, **180**, 211–224.
- Steinkamp, J. & Hickler, T. (2015) Is drought-induced forest dieback globally increasing? *Journal of Ecology*, **103**, 31–42.
- Stocker, T.F., Qin, D., Plattner, G.-K., Alexander, L.V., Allen, S.K., Bindoff, N.L. *et al.* (2013). Technical summary. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds T.F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex & P.M. Midgley), pp. 33–115. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- Vayreda, J., Martínez-Vilalta, J., Gracia, M. & Retana, J. (2012) Recent climate changes interact with stand structure and management to determine changes in tree carbon stocks in Spanish forests. *Global Change Biology*, **18**, 1028–1041.
- Warren, J.M., Norby, R.J. & Wullschleger, S.D. (2011) Elevated CO₂ enhances leaf senescence during extreme drought in a temperate forest. *Tree Physiology*, **31**, 117–130.
- Warszawski, L., Friend, A.D., Ostberg, S., Frieler, K., Lucht, W., Schaphoff, S. *et al.* (2013) A multi-model analysis of risk of ecosystem shifts under climate change. *Environmental Research Letters*, **8**, 044018.
- White, A., Cannell, M.G.R. & Friend, A.D. (1999) Climate change impacts on ecosystems and the terrestrial carbon sink: a new assessment. *Global Environmental Change*, **9**(Suppl. 1), S21–S30.
- Wiens, J.A. (1989) Spatial scaling in ecology. *Functional Ecology*, **3**, 385–397.
- Zaehle, S., Medlyn, B.E., De Kauwe, M.G., Walker, A.P., Dietze, M.C., Hickler, T. *et al.* (2014) Evaluation of 11 terrestrial carbon–nitrogen cycle models against observations from two temperate free-air CO₂ enrichment studies. *New Phytologist*, **202**, 803–822.

Received 22 April 2014; accepted 13 October 2014

Handling Editor: David Coomes